


Føtal påvirkning af hormonforstyrrende stoffer



Margit Bistrup Fisher, Læge, Ph.d.-studerende
Anders Juul, Ledende overlæge, Professor
Casper Hagen, Afdelingslæge

Baggrund

CONSENSUS STATEMENT

OPEN


 Check for updates

Paracetamol use during pregnancy — a call for precautionary action

Ann Z. Bauer¹, Shanna H. Swan², David Kriebel¹, Zeyan Liew³, Hugh S. Taylor⁴, Carl-Gustaf Bornehag^{2,5}, Anderson M. Andrade⁶, Jørn Olsen⁷, Rigmor H. Jensen⁸, Rod T. Mitchell⁹, Niels E. Skakkebaek¹⁰, Bernard Jégou^{11,13} and David M. Kristensen^{8,11,12}


Abstract | Paracetamol (N-acetyl-p-aminophenol (APAP), otherwise known as acetaminophen) is the active ingredient in more than 600 medications used to relieve mild to moderate pain and reduce fever. APAP is widely used by pregnant women as governmental agencies, including the FDA and EMA, have long considered APAP appropriate for use during pregnancy when used as directed. However, increasing experimental and epidemiological research suggests that prenatal exposure to APAP might alter fetal development, which could increase the risks of some neurodevelopmental, reproductive and urogenital disorders. Here we summarize this evidence and call for precautionary action through a focused research effort and by increasing awareness among health professionals and pregnant women. APAP is an important medication

I graviditeten



56%


1-2 x ugt. i 1.trimester



Ca. 235 gravide

~7%

Dagligt i 1.trimester




Ca. 60 gravide




>1%

The Copenhagen Analgesic Study - COPANA

- Formålet med dette studie er at afklare, hvorvidt indtag af mild smertestillende medicin i graviditeten, påvirker udviklingen af æggestokke eller testikler hos fosteret



Føtal AGD



Børneundersøgelsen

Vægt/længde/hudfolder



AGD

Pubertetstegn





Penis-mål

UL testikler / UL uterus og ovarier

Blod- og urinprøve




Ultralyd




COPANA - biobank

3000 urinprøver!



2400 blodprøver!



Graviditetsuge
8-10

Graviditetsuge
12

Graviditetsuge
30-33

Graviditetsuge
37-42

3 måneder
gammel

Web-baseret
spørgeskema

Nakkefoldskanning
Møde med læge

3. trimester ultralydskanning
og mål af anogenitale
afstand

Fødsel

Børneundersøgelse

Tak for jeres opmærksomhed!

Tak til alle familierne!

CeHoS



Aase og Ejnar Danielsens Fond



Læge Sofus Carl Emil Friis og
Hustru Olga Doris Friis' Legat

Axel Muusfeldts Fond

